

## A TWO-PARAMETER EXTENSION OF URBANIK'S PRODUCT CONVOLUTION SEMIGROUP

BY

CHRISTIAN BERG (COPENHAGEN)

*Abstract.* We prove that  $s_n(a, b) = \Gamma(an + b)/\Gamma(b)$ ,  $n = 0, 1, \dots$ , is an infinitely divisible Stieltjes moment sequence for arbitrary  $a, b > 0$ . Its powers  $s_n(a, b)^c$ ,  $c > 0$ , are Stieltjes determinate if and only if  $ac \leq 2$ . The latter was conjectured in a paper by Lin (2019) in the case  $b = 1$ . We describe a product convolution semigroup  $\tau_c(a, b)$ ,  $c > 0$ , of probability measures on the positive half-line with densities  $e_c(a, b)$  and having the moments  $s_n(a, b)^c$ . We determine the asymptotic behavior of  $e_c(a, b)(t)$  for  $t \rightarrow 0$  and for  $t \rightarrow \infty$ , and the latter implies the Stieltjes indeterminacy when  $ac > 2$ . The results extend the previous work of the author and López (2015) and lead to a convolution semigroup of probability densities  $(g_c(a, b)(x))_{c>0}$  on the real line. The special case  $(g_c(a, 1)(x))_{c>0}$  are the convolution roots of the Gumbel distribution with scale parameter  $a > 0$ . All the densities  $g_c(a, b)(x)$  lead to determinate Hamburger moment problems.

**2010 AMS Mathematics Subject Classification:** Primary: 60E07;  
Secondary: 60B15, 44A60.

**Key words and phrases:** Infinitely divisible Stieltjes moment sequence, product convolution semigroup, asymptotic approximation of integrals, Gumbel distribution.

### 1. INTRODUCTION

A Stieltjes moment sequence is a sequence of non-negative numbers of the form

$$(1.1) \quad s_n = \int_0^\infty t^n d\mu(t), \quad n \in \mathbb{N}_0 := \{0, 1, 2, \dots\},$$

where  $\mu$  is a positive measure on  $[0, \infty)$  such that  $t^n \in L^1(\mu)$  for all  $n \in \mathbb{N}_0$ . The sequence  $(s_n)$  is called *normalized* if  $s_0 = \mu([0, \infty)) = 1$ , and it is called *S-determinate* (resp. *S-indeterminate*) if (1.1) has exactly one solution (resp. several solutions)  $\mu$  as positive measures on  $[0, \infty)$ . All these concepts go back to the fundamental memoir of Stieltjes [19].

A Stieltjes moment sequence  $(s_n)$  is called *infinitely divisible* if  $(s_n^c)$  is a Stieltjes moment sequence for any  $c > 0$ . These sequences were characterized in Tyan's Ph.D. thesis [21] and again in [5] without the knowledge of [21]. An important example of an infinitely divisible normalized Stieltjes moment sequence is  $s_n = n!$ , first established in Urbanik [22]. He proved that  $e_c$  in (1.2) is a probability density such that

$$(1.2) \quad (n!)^c = \int_0^\infty t^n e_c(t) dt, \quad e_c(t) = \frac{1}{2\pi} \int_{-\infty}^\infty t^{ix-1} \Gamma(1-ix)^c dx, \quad c, t > 0.$$

Here  $\Gamma$  is Euler's gamma function. The family  $(\tau_c)_{c>0}$  with  $d\tau_c(t) = e_c(t)dt$  is a convolution semigroup in the sense of [6] on the locally compact abelian group  $G = (0, \infty)$  under multiplication. It is called the *Urbanik semigroup* in [7]. It turns out that the terminology "Urbanik semigroup" has been used in the literature for certain semigroups of operators on Banach spaces with the precise name "Urbanik decomposability semigroup", see Section 2 in [10]. We have therefore decided to use the more precise name "Urbanik's product convolution semigroup".

By Carleman's criterion for S-determinacy it is easy to prove that  $(n!)^c$  is S-determinate for  $c \leq 2$ . That this estimate is sharp was first proved in [4], where it was established that  $(n!)^c$  is S-indeterminate for  $c > 2$  based on asymptotic results of Skorokhod [18] about stable distributions, see [23]. Another proof of the S-indeterminacy was given in [7] based on the asymptotic behavior of  $e_c(t)$ ,

$$(1.3) \quad e_c(t) = \frac{(2\pi)^{(c-1)/2} \exp(-ct^{1/c})}{\sqrt{c}} \frac{1}{t^{(c-1)/(2c)}} [1 + O(t^{-1/c})], \quad t \rightarrow \infty.$$

In the recent paper [11], Lin proposes the following conjecture:

CONJECTURE. *Let  $a > 0$  be a real constant and let  $s_n = \Gamma(na + 1)$ ,  $n \in \mathbb{N}_0$ .*

*Then:*

- (a)  *$(s_n)$  is an infinitely divisible Stieltjes moment sequence.*
- (b) *For real  $c > 0$  the sequence  $(s_n^c)$  is S-determinate if and only if  $ac \leq 2$ .*
- (c) *For  $0 < c \leq 2/a$  the unique probability measure  $\mu_c$  corresponding to  $(s_n^c)$*

*has the Mellin transform*

$$\int_0^\infty t^s d\mu_c(t) = \Gamma(as + 1)^c, \quad s \geq 0.$$

When  $a = 1$ , the conjecture is true because of the known results about Urbanik's product convolution semigroup, and for  $a \in \mathbb{N}$ ,  $a \geq 2$ , the conjecture is true because of Theorems 4 and 7 in [11].

We shall prove that the conjecture is true, and it is a special case of similar results for the following more general normalized Stieltjes moment sequence

$$(1.4) \quad s_n(a, b) = \frac{\Gamma(an + b)}{\Gamma(b)} = \frac{1}{a\Gamma(b)} \int_0^\infty t^n t^{b/a-1} \exp(-t^{1/a}) dt, \quad n = 0, 1, \dots,$$

where  $a, b > 0$  are arbitrary.

Defining

$$(1.5) \quad e_1(a, b)(t) = \frac{1}{a\Gamma(b)} t^{b/a-1} \exp(-t^{1/a}),$$

we get for  $\operatorname{Re} z > -b/a$  and after a change of variable  $t = s^a$

$$(1.6) \quad \int_0^\infty t^z e_1(a, b)(t) dt = \Gamma(az + b)/\Gamma(b).$$

This leads to our first main result.

**THEOREM 1.1.** (i)  $(s_n(a, b))$  is an infinitely divisible Stieltjes moment sequence.

(ii) There exists a uniquely determined convolution semigroup  $(\tau_c(a, b))_{c>0}$  of probability measures on the multiplicative group  $(0, \infty)$  such that

$$(1.7) \quad \int_0^\infty t^z d\tau_c(a, b)(t) = [\Gamma(az + b)/\Gamma(b)]^c, \quad \operatorname{Re} z > -b/a,$$

and, in particular,  $(s_n(a, b)^c)$  is the moment sequence of  $\tau_c(a, b)$ .

(iii)  $d\tau_c(a, b)(t) = e_c(a, b)(t) dt$  on  $(0, \infty)$ , where

$$(1.8) \quad e_c(a, b)(t) = \frac{1}{2\pi} \int_{-\infty}^\infty t^{ix-1} [\Gamma(b - iax)/\Gamma(b)]^c dx, \quad t > 0,$$

is a probability density belonging to  $C^\infty(0, \infty)$ .

(iv)  $(s_n(a, b)^c)$  is  $S$ -determinate if and only if  $ac \leq 2$ , hence independent of  $b > 0$ .

Note that (1.4) is a special case of (1.6).

The measure  $\tau_1(a, b)$  was considered in [20], where it was proved that the measure is  $S$ -indeterminate if  $a > \max(2, 2b)$ . This is a consequence of our result. Note that  $\tau_1(a, 1)$  is called the *Weibull distribution* with shape parameter  $1/a$  and scale parameter one.

In (1.7) and (1.8) we use that  $\Gamma(z)$  is a non-vanishing holomorphic function in the cut plane

$$(1.9) \quad \mathcal{A} = \mathbb{C} \setminus (-\infty, 0],$$

so we can define

$$\Gamma(z)^c := \exp(c \log \Gamma(z)), \quad z \in \mathcal{A},$$

using the holomorphic branch of  $\log \Gamma$  which is zero for  $z = 1$ . This branch is explicitly given in (3.1) below.

Let us recall a few facts about convolution semigroups of probability measures on LCA groups, see [6] for details.

The continuous characters of the multiplicative group  $G = (0, \infty)$  can be given as  $t \rightarrow t^{ix}$ , where  $x \in \mathbb{R}$  is arbitrary, and in this way the dual group  $\widehat{G}$  of  $G$  can be identified with the additive group of real numbers. The convolution between measures  $\mu$  and  $\sigma$  on  $(0, \infty)$ , called a *product convolution* and denoted by  $\mu \diamond \sigma$ , is defined as

$$\int_0^\infty f(t) d\mu \diamond \sigma(t) = \int_0^\infty \int_0^\infty f(ts) d\mu(t) d\sigma(s)$$

for suitable classes of continuous functions  $f$  on  $(0, \infty)$ , e.g. those of compact support.

A family  $(\mu_c)_{c>0}$  of probability measures on the multiplicative group  $G = (0, \infty)$  is called a *convolution semigroup* if

$$\mu_c \diamond \mu_d = \mu_{c+d}, \quad c, d > 0, \quad \text{and} \quad \lim_{c \rightarrow 0} \mu_c = \varepsilon_1 \text{ vaguely.}$$

Here  $\varepsilon_1$  is the Dirac measure with total mass one concentrated in the neutral element one of the group. Given a convolution semigroup  $(\mu_c)_{c>0}$  on  $(0, \infty)$ , it is easy to see that if  $\mu_1$  has moments of order  $n$ , then all the measures  $\mu_c$  have moments of order  $n$  and

$$\int_0^\infty t^n d\mu_c(t) = \left( \int_0^\infty t^n d\mu_1(t) \right)^c, \quad c > 0.$$

By [6], Theorem 8.3, there is a one-to-one correspondence between convolution semigroups  $(\mu_c)_{c>0}$  of probability measures on  $G$  and continuous negative definite functions  $\rho : \mathbb{R} \rightarrow \mathbb{C}$  satisfying  $\rho(0) = 0$  such that

$$(1.10) \quad \int_0^\infty t^{-ix} d\mu_c(t) = \exp(-c\rho(x)), \quad c > 0, x \in \mathbb{R}.$$

By the inversion theorem of Fourier analysis for LCA groups, if  $\exp(-c\rho)$  is integrable on  $\mathbb{R}$ , then  $d\mu_c(t) = f_c(t) dt$  for a continuous function  $f_c(t)$  ( $tf_c(t)$  is the density of  $\mu_c$  with respect to Haar measure  $(1/t)dt$  on  $(0, \infty)$ ) given by

$$(1.11) \quad f_c(t) = \frac{1}{2\pi} \int_{-\infty}^\infty t^{ix-1} \exp(-c\rho(x)) dx, \quad t > 0.$$

(Note that the dual Haar measure of  $(1/t)dt$  on  $(0, \infty)$  is  $1/(2\pi) dx$  on  $\mathbb{R}$ .)

PROPOSITION 1.1. For  $a, b > 0$

$$(1.12) \quad \rho(x) := \log \Gamma(b) - \log \Gamma(b - iax), \quad x \in \mathbb{R},$$

is a continuous negative definite function on  $\mathbb{R}$  satisfying  $\rho(0) = 0$ .

Proposition 1.1 shows that there exists a uniquely determined product convolution semigroup  $(\tau_c(a, b))_{c>0}$  satisfying

$$(1.13) \quad \int_0^\infty t^{-ix} d\tau_c(a, b)(t) = \exp[-c(\log \Gamma(b) - \log \Gamma(b - iax))] \\ = [\Gamma(b - iax)/\Gamma(b)]^c, \quad x \in \mathbb{R}.$$

Putting  $z = -ix$  in (1.6), we see by the uniqueness theorem for Fourier transforms that  $d\tau_1(a, b)(t) = e_1(a, b)(t) dt$ , and since  $e_1(a, b)(t)$  has moments of any order by (1.6), we infer that all the measures  $\tau_c(a, b)$  have moments of any order. This implies that the integral

$$\int_0^\infty t^z d\tau_c(a, b)(t), \quad \operatorname{Re} z \geq 0,$$

defines a continuous function of  $z$  in the half-plane  $\operatorname{Re} z \geq 0$  and holomorphic in the interior  $\operatorname{Re} z > 0$ . By (1.13) this function equals  $[\Gamma(b + az)/\Gamma(b)]^c$  on the imaginary axis and hence on  $\operatorname{Re} z \geq 0$ . As in the proof of [4], Lemma 2.1, it follows that this equality extends to the half-plane  $\operatorname{Re} z > -b/a$ , i.e. (1.7) holds.

The function  $(\Gamma(b - iax)/\Gamma(b))^c$  is a Schwartz function on  $\mathbb{R}$  and in particular integrable, so (1.8) follows from (1.7), and  $e_c(a, b)$  is  $C^\infty$  on  $(0, \infty)$ .

In this way we have established (i)–(iii) of Theorem 1.1. The proof of the more difficult part (iv) as well as the proof of Proposition 1.1 will be given in Section 3.

By Riemann–Lebesgue's lemma we also see that  $te_c(a, b)(t)$  tends to zero for  $t$  tending to zero and to infinity. Much more on the behavior near zero and infinity will be given in Section 2, where we extend the work of [7] leading to the asymptotic behavior of the densities  $e_c(a, b)(t)$  for  $t \rightarrow 0$  and  $t \rightarrow \infty$ . The behavior for  $t \rightarrow \infty$  will lead to a proof of the S-indeterminacy for  $ac > 2$  using the Krein criterion.

The fact that  $\tau_c(a, b) \diamond \tau_d(a, b) = \tau_{c+d}(a, b)$  can be written as

$$(1.14) \quad e_{c+d}(a, b)(t) = \int_0^\infty e_c(a, b)(t/x) e_d(a, b)(x) \frac{dx}{x}, \quad c, d > 0.$$

In particular, for  $c = d = 1$  and the explicit formula for  $e_1(a, b)$  we get

$$(1.15) \quad e_2(a, b)(t) = \frac{t^{b/a-1}}{[a\Gamma(b)]^2} \int_0^\infty \exp(-x^{-1/a} t^{1/a} - x^{1/a}) \frac{dx}{x} \\ = \frac{2t^{b/a-1}}{a\Gamma(b)^2} K_0(2t^{1/(2a)}),$$

because the Macdonald function  $K_0$  is given by

$$K_0(z) = \frac{1}{2} \int_0^\infty \exp\left(-\frac{(z/2)^2}{y-y}\right) \frac{dy}{y},$$

cf. [8], 8.432(6); [14], Chapter 10, Section 25.

Except for a scaling this result is the same as Lemma 1 in [12].

## 2. MAIN RESULTS

Our additional main results are the following:

**THEOREM 2.1.** *For  $c > 0$  we have*

$$(2.1) \quad e_c(a, b)(t) = \frac{(2\pi)^{(c-1)/2} \exp(-ct^{1/(ac)})}{a\sqrt{c}\Gamma(b)^c t^{1-(b-1/2+1/(2c))/a}} [1 + \mathcal{O}(t^{-1/(ac)})], \quad t \rightarrow \infty.$$

**THEOREM 2.2.** *The measure  $\tau_c(a, b)$  is  $S$ -indeterminate if and only if  $ac > 2$ .*

**THEOREM 2.3.** *For  $c > 0$  and  $0 < t < 1$  we have*

$$(2.2) \quad e_c(a, b)(t) = \frac{t^{b/a-1} [\log(1/t)]^{c-1}}{[a\Gamma(b)]^c \Gamma(c)} + \mathcal{O}(t^{b/a-1} [\log(1/t)]^{c-2}), \quad t \rightarrow 0.$$

**REMARK 2.1.** Formula (2.2) shows that  $e_c(a, b)(t)$  tends to zero for  $t \rightarrow 0$  if  $b/a > 1$ , and to infinity if  $b/a < 1$ , independent of  $c$ . If  $b/a = 1$ , then  $e_c(a, b)(t)$  tends to zero for  $c < 1$  and to infinity as a power of  $\log(1/t)$  when  $c > 1$ .

## 3. PROOFS

**Proof of Proposition 1.1.** From the Weierstrass product for the entire function  $1/\Gamma(z)$  we get the following holomorphic branch in the cut plane  $\mathcal{A}$ , cf. (1.9):

$$(3.1) \quad -\log \Gamma(z) = \gamma z + \text{Log } z + \sum_{k=1}^{\infty} (\text{Log}(1 + z/k) - z/k), \quad z \in \mathcal{A},$$

where  $\text{Log}$  denotes the principal logarithm, and  $\gamma$  is Euler's constant.

For  $n \in \mathbb{N}$  and  $z \in \mathcal{A}$  define

$$\rho_n(z) = \gamma z + \text{Log } z + \sum_{k=1}^n (\text{Log}(1 + z/k) - z/k),$$

$$R_n(z) = \sum_{k=n+1}^{\infty} (\text{Log}(1 + z/k) - z/k),$$

so  $\lim_{n \rightarrow \infty} \rho_n(z) = -\log \Gamma(z)$ , uniformly on compact subsets of  $\mathcal{A}$ . Furthermore, we have

$$\log \Gamma(b) + \rho_n(b) + R_n(b) = 0,$$

and since  $\log(1+x) < x$  for  $x > 0$ , we see that  $R_n(b) < 0$  and hence  $\log \Gamma(b) + \rho_n(b) > 0$ .

We claim that  $\log \Gamma(b) + \rho_n(b - iax)$  is a continuous negative definite function, and letting  $n \rightarrow \infty$ , we get the assertion of Proposition 1.1.

To see the claim, we write

$$\begin{aligned} \log \Gamma(b) + \rho_n(b - iax) &= \log \Gamma(b) + (b - iax) \left( \gamma - \sum_{k=1}^n \frac{1}{k} \right) \\ &\quad + \text{Log}(b - iax) + \sum_{k=1}^n \text{Log} \left( 1 + \frac{b - iax}{k} \right) \\ &= \log \Gamma(b) + \rho_n(b) - iax \left( \gamma - \sum_{k=1}^n \frac{1}{k} \right) + \sum_{k=0}^n \text{Log} \left( 1 - i \frac{ax}{b+k} \right), \end{aligned}$$

and the assertion follows since  $\alpha + i\beta x$  and  $\text{Log}(1 + i\beta x)$  are negative definite functions when  $\alpha \geq 0, \beta \in \mathbb{R}$ , see [6], [16]. ■

**Proof of Theorem 2.1.** We modify the proof given in [7] and start by applying Cauchy's integral theorem to move the integration in (1.8) to a horizontal line

$$(3.2) \quad H_\delta := \{z = x + i\delta : x \in \mathbb{R}\}, \quad \delta > -b/a.$$

LEMMA 3.1. *With  $H_\delta$  as in (3.2) we have*

$$(3.3) \quad e_c(a, b)(t) = \frac{1}{2\pi} \int_{H_\delta} t^{iz-1} [\Gamma(b - iaz)/\Gamma(b)]^c dz, \quad t > 0.$$

**Proof.** For  $t, c > 0$  fixed,  $f(z) = t^{iz-1} [\Gamma(b - iaz)/\Gamma(b)]^c$  is holomorphic in the simply connected domain  $\mathbb{C} \setminus i(-\infty, -b/a]$ , so (3.3) follows from Cauchy's integral theorem provided the integral

$$\int_0^\delta f(x + iy) dy$$

tends to zero for  $x \rightarrow \pm\infty$ . We have

$$|f(x + iy)| = t^{-y-1} |\Gamma(b + y - iax)/\Gamma(b)|^c,$$

and the result follows since

$$|\Gamma(u + iv)| \sim \sqrt{2\pi} e^{-\pi|v|/2} |v|^{u-1/2}, \quad |v| \rightarrow \infty, \quad \text{uniformly for bounded real } u,$$

cf. [1], p. 141, equation 5.11.9; [8], 8.328(1). ■

In the following we will use Lemma 3.1 with the line of integration  $H_\delta$ , where  $\delta = (t^{1/(ac)} - b)/a$ . Therefore,

$$e_c(a, b)(t) = t^{(b-t^{1/(ac)})/a-1} \frac{1}{2\pi} \int_{-\infty}^{\infty} t^{ix} [\Gamma(t^{1/(ac)} - iax)/\Gamma(b)]^c dx,$$

and after the change of variable  $x = a^{-1}t^{1/(ac)}u$  and putting  $A := (1/c + b - a)/a$  we get

(3.4)

$$e_c(a, b)(t) = t^{A-a^{-1}t^{1/(ac)}} \frac{1}{2\pi a} \int_{-\infty}^{\infty} t^{iua^{-1}t^{1/(ac)}} [\Gamma(t^{1/(ac)}(1 - iu))/\Gamma(b)]^c du.$$

Binet's formula for  $\Gamma$  is ([8], 8.341(1))

$$(3.5) \quad \Gamma(z) = \sqrt{2\pi} z^{z-1/2} e^{-z+\mu(z)}, \quad \operatorname{Re}(z) > 0,$$

where

$$(3.6) \quad \mu(z) = \int_0^\infty \left( \frac{1}{2} - \frac{1}{t} + \frac{1}{e^t - 1} \right) \frac{e^{-zt}}{t} dt, \quad \operatorname{Re}(z) > 0.$$

Notice that  $\mu(z)$  is the Laplace transform of a positive function, so we have the estimates for  $z = r + is, r > 0$ , that is,

$$(3.7) \quad |\mu(z)| \leq \mu(r) \leq \frac{1}{12r},$$

where the last inequality is a classical version of Stirling's formula, thus showing that the estimate is uniform in  $s \in \mathbb{R}$ .

Inserting this in (3.4), we get after some simplification

(3.8)

$$e_c(a, b)(t) = \frac{(2\pi)^{c/2-1}}{a\Gamma(b)^c} t^{A-1/(2a)} e^{-ct^{1/(ac)}} \int_{-\infty}^{\infty} e^{ct^{1/(ac)}f(u)} g_c(u) M(u, t) du,$$

where

$$(3.9) \quad f(u) := iu + (1 - iu) \operatorname{Log}(1 - iu), \quad g_c(u) := (1 - iu)^{-c/2}$$

and

$$(3.10) \quad M(u, t) := \exp [c\mu(t^{1/(ac)}(1 - iu))].$$

From (3.7) we get  $M(u, t) = 1 + \mathcal{O}(t^{-1/(ac)})$  for  $t \rightarrow \infty$ , uniformly in  $u$ . We shall therefore consider the behavior for large  $x$  of the integral

$$(3.11) \quad \int_{-\infty}^{\infty} e^{xf(u)} g_c(u) du, \quad x = ct^{1/(ac)}.$$



This is the same integral as was treated in [7], equation (28), leading to

$$\int_{-\infty}^{\infty} e^{xf(u)} g_c(u) du = (2\pi/x)^{1/2} [1 + \mathcal{O}(x^{-1})]$$

by methods from [13].

For  $x = ct^{1/(ac)}$  we find

$$\int_{-\infty}^{\infty} \exp(ct^{1/(ac)} f(u)) g_c(u) du = \frac{\sqrt{2\pi}}{\sqrt{ct^{1/(2ac)}}} [1 + \mathcal{O}(t^{-1/(ac)})],$$

hence

$$e_c(a, b)(t) = \frac{(2\pi)^{(c-1)/2}}{a\sqrt{c}\Gamma(b)^c} \frac{\exp(-ct^{1/(ac)})}{t^{1-(b-1/2+1/(2c))/a}} [1 + \mathcal{O}(t^{-1/(ac)})]. \quad \blacksquare$$

**Proof of Theorem 2.2.** We first prove that  $(s_n(a, b)^c)$  is S-determinate for  $ac \leq 2$  by Carleman's criterion, cf. [17], p. 20. In fact, from Stirling's formula we have

$$s_n(a, b)^{c/(2n)} = (\Gamma(na + b)/\Gamma(b))^{c/(2n)} \sim (na/e)^{ac/2}, \quad n \rightarrow \infty,$$

so  $\sum s_n(a, b)^{-c/(2n)} = \infty$  if and only if  $ac \leq 2$ .

Since Carleman's criterion is only a sufficient condition for S-determinacy, we need to prove that  $e_c(a, b)$  is S-indeterminate for  $ac > 2$ . We apply the Krein criterion for S-indeterminacy of probability densities concentrated on the half-line, using a version due to H. L. Pedersen given in [9], Theorem 4. It states that if

$$(3.12) \quad \int_K^{\infty} \frac{\log e_c(a, b)(t^2) dt}{1 + t^2} > -\infty$$

for some  $K \geq 0$ , then  $e_c(a, b)$  is S-indeterminate. This version of the Krein criterion is a simplification of a stronger version given in [15]. We shall see that (3.12) holds for  $ac > 2$ .

From Theorem 2.1 we see that (3.12) holds for sufficiently large  $K > 0$  if and only if

$$\int_K^{\infty} \frac{-ct^{2/(ac)}}{1 + t^2} dt > -\infty,$$

and the latter holds precisely for  $ac > 2$ . This shows that  $\tau_c(a, b)$  is S-indeterminate for  $ac > 2$ .  $\blacksquare$

**Proof of Theorem 2.3.** The proof uses the same ideas as in [7], but since the proof is quite technical, we give the full proof with the necessary modifications. Since we are studying the behavior for  $t \rightarrow 0$ , we assume that  $0 < t < 1$  so that  $\Lambda := \log(1/t) > 0$ .

We will need integration along vertical lines

$$(3.13) \quad V_\alpha := \{\alpha + iy \mid y = -\infty \dots \infty\}, \quad \alpha \in \mathbb{R},$$

and we can therefore express (1.8) as

$$(3.14) \quad e_c(a, b)(t) = \frac{t^{b/a-1}}{2\pi ia\Gamma(b)^c} \int_{V_{-b}} t^{z/a}\Gamma(-z)^c dz.$$

By the functional equation for  $\Gamma$  we get

$$(3.15) \quad e_c(a, b)(t) = \frac{t^{b/a-1}}{2\pi ia\Gamma(b)^c} \int_{V_{-b}} g(z)\varphi(z) dz,$$

where we have defined

$$\varphi(z) := t^{z/a}\Gamma(1-z)^c, \quad g(z) := (-z)^{-c} = \exp(-c \operatorname{Log}(-z)).$$

Note that  $\varphi$  is holomorphic in  $\mathbb{C} \setminus [1, \infty)$ , while  $g$  is holomorphic in  $\mathbb{C} \setminus [0, \infty)$ .

For  $x > 0$  we define

$$g_\pm(x) := \lim_{\varepsilon \rightarrow 0^+} g(x \pm i\varepsilon) = x^{-c} e^{\pm i\pi c}.$$

**Case 1.** Assume  $0 < c < 1$ .

We fix  $0 < s < 1$ , choose  $0 < \varepsilon < \min(s, b)$  and integrate  $g(z)\varphi(z)$  over the contour

$$\begin{aligned} \mathcal{C} := & \{-b + iy \mid y = \infty \dots 0\} \cup [-b, -\varepsilon] \cup \{\varepsilon e^{i\theta} \mid \theta = \pi \dots 0\} \\ & \cup [\varepsilon, s] \cup \{s + iy \mid y = 0 \dots \infty\} \end{aligned}$$

and get zero by the integral theorem of Cauchy. On the interval  $[\varepsilon, s]$  we will use  $g = g_+$ .

Similarly, we get zero by integrating  $g(z)\varphi(z)$  over the complex conjugate contour  $\bar{\mathcal{C}}$ , and now we use  $g = g_-$  on the interval  $[\varepsilon, s]$ .

Subtracting the second contour integral from the first leads to

$$\int_{V_s} - \int_{V_{-b}} - \int_{|z|=\varepsilon} g(z)\varphi(z) dz + \int_{\varepsilon}^s \varphi(x)(g_+(x) - g_-(x)) dx = 0,$$

where the integral over the circle is with positive orientation. Note that the two integrals over  $[-b, -\varepsilon]$  cancel. Since  $0 < c < 1$ , it is easy to see that the just mentioned integral converges to zero for  $\varepsilon \rightarrow 0$ , and we finally get for  $\varepsilon \rightarrow 0$

$$\begin{aligned} e_c(a, b)(t) &= \frac{t^{b/a-1}}{2\pi ia\Gamma(b)^c} \int_{V_s} g(z)\varphi(z) dz + \frac{t^{b/a-1} \sin(\pi c)}{\pi a\Gamma(b)^c} \int_0^s x^{-c}\varphi(x) dx \\ &:= I_1 + I_2. \end{aligned}$$

We claim that  $I_1$  is  $o(t^{(s+b)/a-1})$  for  $t \rightarrow 0$ . To see this, we insert the parametrization of  $V_s$  and get

$$\begin{aligned} I_1 &= \frac{t^{b/a-1}}{2\pi a\Gamma(b)^c} \int_{-\infty}^{\infty} (-s - iy)^{-c} t^{(s+iy)/a} \Gamma(1 - s - iy)^c dy \\ &= \frac{t^{(s+b)/a-1}}{2\pi a\Gamma(b)^c} \int_{-\infty}^{\infty} e^{-iy\Lambda/a} (-s - iy)^{-c} \Gamma(1 - s - iy)^c dy, \end{aligned}$$

and the integral is  $o(1)$  for  $t \rightarrow 0$  by Riemann–Lebesgue’s lemma because  $\Lambda := \log(1/t) \rightarrow \infty$ .

The substitution  $u = x\Lambda$  in the integral in the term  $I_2$  leads to

$$(3.16) \quad I_2 = \frac{t^{b/a-1} \sin(\pi c)}{\pi a\Gamma(b)^c} \Lambda^{c-1} \int_0^{s\Lambda} u^{-c} e^{-u/a} \Gamma(1 - u/\Lambda)^c du.$$

We split the integral in (3.16) as

$$(3.17) \quad \int_0^{s\Lambda} u^{-c} e^{-u/a} [\Gamma(1 - u/\Lambda)^c - 1] du + \int_0^{\infty} u^{-c} e^{-u/a} du - \int_{s\Lambda}^{\infty} u^{-c} e^{-u/a} du.$$

Calling the three terms  $J_1, J_2, J_3$ , we have  $J_2 = a^{1-c}\Gamma(1 - c)$  and

$$J_3 = -a^{1-c}\Gamma(1 - c, s\Lambda/a),$$

where  $\Gamma(\alpha, x)$  is the incomplete gamma function with the asymptotics

$$\Gamma(\alpha, x) = \int_x^{\infty} u^{\alpha-1} e^{-u} du \sim x^{\alpha-1} e^{-x}, \quad x \rightarrow \infty$$

(cf. [8], 8.357), hence

$$J_3 = \mathcal{O}(t^{s/a}\Lambda^{-c}), \quad t \rightarrow 0.$$

Using the digamma function  $\Psi = \Gamma'/\Gamma$ , we get by the mean-value theorem

$$\Gamma(1 - u/\Lambda)^c - 1 = -\frac{u}{\Lambda} c\Gamma(1 - \theta u/\Lambda)^c \Psi(1 - \theta u/\Lambda)$$

for some  $0 < \theta < 1$ , but this implies that

$$|\Gamma(1 - u/\Lambda)^c - 1| \leq \frac{cu}{\Lambda} M(s), \quad 0 < u < s\Lambda,$$

where

$$M(s) := \max\{\Gamma(x)^c |\Psi(x)| \mid 1 - s \leq x \leq 1\},$$

so  $J_1 = \mathcal{O}(\Lambda^{-1})$  for  $t \rightarrow 0$ .

This gives

$$\begin{aligned} I_2 &= \frac{t^{b/a-1} \sin(\pi c)}{\pi a \Gamma(b)^c} \Lambda^{c-1} (\mathcal{O}(\Lambda^{-1}) + a^{1-c} \Gamma(1-c) + \mathcal{O}(t^{s/a} \Lambda^{-c})) \\ &= \frac{t^{b/a-1} \Lambda^{c-1}}{(a \Gamma(b))^c \Gamma(c)} + \mathcal{O}(t^{b/a-1} \Lambda^{c-2}), \end{aligned}$$

where we have used Euler's reflection formula for  $\Gamma$ . Since finally

$$I_1 = o(t^{(s+b)/a-1}) = \mathcal{O}(t^{b/a-1} \Lambda^{c-2}),$$

we see that (2.2) holds.

**C a s e 2.** Assume  $1 < c < 2$ .

The gamma function decays so rapidly on vertical lines  $z = \alpha + iy$ ,  $y \rightarrow \pm\infty$ , that we can integrate by parts in (3.15) to get

$$(3.18) \quad e_c(a, b)(t) = -\frac{t^{b/a-1}}{2\pi i a \Gamma(b)^c} \int_{V_{-1}} \frac{(-z)^{-(c-1)}}{c-1} \frac{d}{dz} (t^{z/a} \Gamma(1-z)^c) dz.$$

Defining

$$\varphi_1(z) := \frac{d}{dz} (t^{z/a} \Gamma(1-z)^c) = t^{z/a} \Gamma(1-z)^c ((1/a) \log t - c \Psi(1-z))$$

and using the same contour technique as in Case 1 to the integral in (3.18), where now  $0 < c-1 < 1$ , we get for  $0 < s < 1$  fixed the equality

$$e_c(a, b)(t) = -\frac{t^{b/a-1}}{a \Gamma(b)^c} (\tilde{I}_1 + \tilde{I}_2),$$

where

$$\tilde{I}_1 = \frac{1}{2\pi i (c-1)} \int_{V_s} (-z)^{-(c-1)} \varphi_1(z) dz, \quad \tilde{I}_2 = \frac{\sin(\pi(c-1))}{\pi(c-1)} \int_0^s x^{-(c-1)} \varphi_1(x) dx.$$

We have  $\tilde{I}_1 = o(t^{s/a} \Lambda)$  for  $t \rightarrow 0$  by Riemann–Lebesgue's lemma, and the substitution  $u = x\Lambda$  in the second integral leads to

$$\begin{aligned} & \int_0^s x^{-(c-1)} \varphi_1(x) dx \\ &= \Lambda^{c-2} \int_0^{s\Lambda} u^{-(c-1)} \varphi_1(u/\Lambda) du \\ &= -(1/a) \Lambda^{c-1} \left( \int_0^{s\Lambda} u^{-(c-1)} e^{-u/a} du + \int_0^{s\Lambda} u^{-(c-1)} e^{-u/a} (\Gamma(1-u/\Lambda)^c - 1) du \right) \end{aligned}$$

$$\begin{aligned} & -c\Lambda^{c-2} \int_0^{s\Lambda} u^{-(c-1)} e^{-u/a} \Gamma(1-u/\Lambda)^c \Psi(1-u/\Lambda) du \\ & = -a^{1-c} \Lambda^{c-1} \Gamma(2-c) + \mathcal{O}(\Lambda^{c-2}). \end{aligned}$$

Using the equality

$$\frac{\sin(\pi(c-1))}{(c-1)\pi} (-a^{1-c} \Lambda^{c-1} \Gamma(2-c)) = -a^{1-c} \frac{\Lambda^{c-1}}{\Gamma(c)}$$

obtained by Euler's reflection formula, we see that (2.2) holds.

**C a s e 3.** Assume  $c > 2$ .

We perform the change of variable  $w = (1/a)\Lambda z$  in (3.15) and assume that  $\Lambda > a$ . This gives

$$e_c(a, b)(t) = \frac{t^{b/a-1} \Lambda^{c-1}}{[a\Gamma(b)]^c} \frac{1}{2\pi i} \int_{V_{-(b/a)\Lambda}} (-w)^{-c} e^{-w} \Gamma(1-aw/\Lambda)^c dw.$$

Using Cauchy's integral theorem, we can shift the contour  $V_{-(b/a)\Lambda}$  to  $V_{-1}$  as the integrand is holomorphic in the vertical strip between both paths and exponentially small at both extremes of that vertical strip. For the holomorphic function  $h(z) = \Gamma(1-z)^c$  in the domain  $G = \mathbb{C} \setminus [1, \infty)$ , which is star-shaped with respect to zero, we have

$$h(z) = h(0) + z \int_0^1 h'(uz) du, \quad z \in G,$$

hence

$$(3.19) \quad \Gamma(1-aw/\Lambda)^c = 1 - \frac{caw}{\Lambda} \int_0^1 \Gamma(1-uaw/\Lambda)^c \Psi(1-uaw/\Lambda) du.$$

Defining

$$R(w) = \int_0^1 \Gamma(1-uaw/\Lambda)^c \Psi(1-uaw/\Lambda) du,$$

we get

$$\begin{aligned} & \frac{1}{2\pi i} \int_{V_{-1}} (-w)^{-c} e^{-w} \Gamma(1-aw/\Lambda)^c dw \\ & = \frac{1}{2\pi i} \int_{V_{-1}} (-w)^{-c} e^{-w} dw + \frac{ac/\Lambda}{2\pi i} \int_{V_{-1}} (-w)^{1-c} e^{-w} R(w) dw. \end{aligned}$$

For any  $w \in V_{-1}$ ,  $0 \leq u \leq 1$  and for  $\Lambda \geq a$  it follows that  $1-uaw/\Lambda$  belongs to the closed vertical strip located between the vertical lines  $V_1$  and  $V_2$ . Because

$\Gamma(z)^c \Psi(z)$  is continuous and bounded in this strip,  $R(w)$  is bounded for  $w \in V_{-1}$  by a constant independent of  $\Lambda \geq a$ . Furthermore,  $(-w)^{1-c} e^{-w}$  is integrable over  $V_{-1}$  because  $c > 2$ .

On the other hand, in the integral

$$\frac{1}{2\pi i} \int_{V_{-1}} (-w)^{-c} e^{-w} dw$$

the contour  $V_{-1}$  may be deformed to a Hankel contour

$$\begin{aligned} \mathcal{H} := & \{x - i \mid x = \infty \dots 0\} \cup \{e^{i\theta} \mid \theta = -\pi/2 \dots -3\pi/2\} \\ & \cup \{x + i \mid x = 0 \dots \infty\} \end{aligned}$$

surrounding  $[0, \infty)$ , and the integral over  $\mathcal{H}$  is Hankel's integral representation of the reciprocal gamma function:

$$\frac{1}{2\pi i} \int_{\mathcal{H}} (-w)^{-c} e^{-w} dw = \frac{1}{\Gamma(c)}.$$

Therefore, when we join everything, we obtain for  $c > 2$ :

$$e_c(a, b)(t) = \frac{t^{b/a-1}}{[a\Gamma(b)]^c} \frac{[\log(1/t)]^{c-1}}{\Gamma(c)} + \mathcal{O}(t^{b/a-1} [\log(1/t)]^{c-2}), \quad t \rightarrow 0.$$

**Case 4.**  $c = 1, c = 2$ .

These cases are easy since  $e_1(a, b)(t)$  is explicitly given by (1.5) and  $e_2(a, b)(t)$  by (1.15). The asymptotics of  $K_0$  is known:

$$K_0(t) = \log(2/t) + \mathcal{O}(1), \quad t \rightarrow 0. \quad \blacksquare$$

**REMARK 3.1.** The behavior of  $e_c(a, b)(t)$  for  $t \rightarrow 0$  can be obtained from (3.14) by using the residue theorem when  $c$  is a natural number. In fact, in this case  $\Gamma(-z)^c$  has a pole of order  $c$  at  $z = 0$ , and a shift of the contour  $V_{-1}$  to  $V_s$ , where  $0 < s < 1$ , has to be compensated by a residue, which will give the behavior for  $t \rightarrow 0$ .

When  $c$  is a natural number, one can actually express  $e_c(a, b)(t)$  in terms of Meijer's  $G$ -function:

$$e_c(a, b)(t) = \frac{t^{b/a-1}}{a\Gamma(b)^c} G_{0,c}^{c,0} \left( t^{1/a} \mid \begin{matrix} - & \cdots & - \\ 0 & \cdots & 0 \end{matrix} \right),$$

cf. Section 9.3 in [8].

## 4. ONE-PARAMETER EXTENSION OF THE GUMBEL DISTRIBUTIONS

The group isomorphism  $x = \log(1/t)$  of the multiplicative group  $(0, \infty)$  onto the additive group  $\mathbb{R}$  transforms the convolution semigroup  $(\tau_c(a, b))_{c>0}$  into an ordinary convolution semigroup  $(G_c(a, b))_{c>0}$  of probability measures on  $\mathbb{R}$  with densities given by

$$(4.1) \quad g_c(a, b)(x) = e^{-x} e_c(a, b)(e^{-x}), \quad x \in \mathbb{R},$$

and  $a, b, c > 0$  are arbitrary. For  $c = 1$  we have

$$(4.2) \quad g_1(a, b)(x) = \frac{1}{a\Gamma(b)} \exp(-bx/a - e^{-x/a}), \quad x \in \mathbb{R}.$$

This density is infinitely divisible and the uniquely determined convolution roots are given by (4.1).

The special density  $g_1(a, 1)(x)$  is the Gumbel density with scale parameter  $a > 0$ , and the basic case  $a = 1$  is discussed in [7]. From the asymptotic behavior of  $e_c(a, b)$  in Theorems 2.1 and 2.3 we can obtain the asymptotic behavior of the convolution roots  $g_c(a, b)$ :

$$(4.3) \quad g_c(a, b)(x) = \frac{(2\pi)^{(c-1)/2}}{a\sqrt{c}\Gamma(b)^c} \frac{\exp(-ce^{-x/(ac)})}{\exp(x(b - 1/2 + 1/(2c))/a)} \left[ 1 + \mathcal{O}\left(\exp(x/(ac))\right) \right]$$

for  $x \rightarrow -\infty$ , and

$$(4.4) \quad g_c(a, b)(x) = \frac{\exp(-bx/a)x^{c-1}}{[a\Gamma(b)]^c\Gamma(c)} + \mathcal{O}\left(\exp(-bx/a)x^{c-2}\right), \quad x \rightarrow \infty.$$

**THEOREM 4.1.** *All densities  $g_c(a, b)$  belong to determinate Hamburger moment problems.*

**Proof.** We first prove that  $g_1(a, b)$  is determinate, and for this it suffices to verify that the moments

$$(4.5) \quad s_n = \int_{-\infty}^{\infty} x^n g_1(a, b)(x) dx$$

satisfy Carleman's condition  $\sum_{n=0}^{\infty} s_{2n}^{-1/(2n)} = \infty$  (cf. [17], p. 19). From (4.5) we get

$$\begin{aligned} s_{2n} &= \frac{1}{a\Gamma(b)} \int_0^{\infty} (\log t)^{2n} t^{b/a-1} \exp(-t^{1/a}) dt = \frac{1}{\Gamma(b)} \int_0^{\infty} (a \log s)^{2n} s^{b-1} e^{-s} ds \\ &< \frac{a^{2n}}{\Gamma(b)} \left( \int_0^1 (\log s)^{2n} s^{b-1} ds + \int_1^{\infty} s^{2n+b-1} e^{-s} ds \right). \end{aligned}$$

By changing variables we see that

$$\int_0^1 (\log s)^{2n} s^{b-1} ds = \frac{(2n)!}{b^{2n+1}},$$

and

$$\int_1^\infty s^{2n+b-1} e^{-s} ds < \Gamma(2n+b),$$

hence

$$s_{2n}^{1/(2n)} < \frac{a}{\Gamma(b)^{1/(2n)}} \left[ \left( \frac{(2n)!}{b^{2n+1}} \right)^{1/(2n)} + \Gamma(2n+b)^{1/(2n)} \right],$$

and the Carleman condition follows from Stirling's formula, which shows that the right-hand side is bounded by  $Kn$  for sufficiently large  $K > 0$ . We next use Corollary 3.3 in [2] to infer that the Carleman condition also holds for all convolution roots  $g_c(a, b)$ . ■

Concerning the moments

$$(4.6) \quad s_n(c) = \int_{-\infty}^{\infty} x^n g_c(a, b)(x) dx, \quad n \in \mathbb{N}_0,$$

of the convolution roots we have the following result:

**THEOREM 4.2.** *The moment  $s_n(c)$  of (4.6) is a polynomial*

$$(4.7) \quad s_n(c) = \sum_{k=1}^n a_{n,k} c^k, \quad n \geq 1,$$

*of degree at most  $n$  in the variable  $c$ . The coefficients  $a_{n,k}$  are given below.*

**Proof.** From (1.7) we get

$$\int_{-\infty}^{\infty} e^{-ixy} dG_c(a, b)(x) = \int_0^{\infty} t^{iy} e_c(a, b)(t) dt = [\Gamma(b+iy)/\Gamma(b)]^c,$$

which shows that the negative definite function  $\rho$  corresponding to the convolution semigroup  $(G_c(a, b))_{c>0}$  is

$$\rho(y) = \log \Gamma(b) - \log \Gamma(b+iy), \quad y \in \mathbb{R}.$$

The derivatives of  $\rho$  can be expressed in terms of the digamma function  $\Psi$ , namely

$$\rho^{(n+1)}(y) = -(ia)^{n+1} \Psi^{(n)}(b+iy), \quad n \in \mathbb{N}_0,$$



so if for  $n \in \mathbb{N}_0$  we define (cf. [3], equation (2.7))

$$\sigma_n := -i^{n+1} \rho^{(n+1)}(0) = (-a)^{n+1} \Psi^{(n)}(b),$$

we find

$$\sigma_0 = a\gamma + \frac{a}{b} - ab \sum_{k=1}^{\infty} \frac{1}{k(b+k)},$$

$$\sigma_n = a^{n+1} n! \sum_{k=0}^{\infty} \frac{1}{(b+k)^{n+1}} = a^{n+1} n! \zeta(n+1, b), \quad n \in \mathbb{N},$$

where  $\zeta(z, q)$  is Hurwitz' zeta function (cf. [8], 9.521).

According to [3] we have  $s_1(c) = \sigma_0 c$ ,  $s_2(c) = \sigma_1 c + \sigma_0^2 c^2$  and in general  $s_n(c)$  is given by (4.7), where the coefficients  $a_{n,k}$  are determined by the recursion

$$a_{n+1,k+1} = \sum_{j=k}^n a_{j,k} \binom{n}{j} \sigma_{n-j}, \quad n \geq k \geq 0.$$

It is easy to see that

$$a_{n,1} = \sigma_{n-1}, \quad a_{n,n-1} = \binom{n}{2} \sigma_0^{n-2} \sigma_1, \quad a_{n,n} = \sigma_0^n. \quad \blacksquare$$

## REFERENCES

- [1] R. A. Askey and R. Roy, *Chapter 5: Gamma Function*, in: *NIST Handbook of Mathematical Functions*, NIST and Cambridge University Press, 2010.
- [2] C. Berg, *On the preservation of determinacy under convolution*, Proc. Amer. Math. Soc. 93 (2) (1985), pp. 351–357.
- [3] C. Berg, *On infinitely divisible solutions to indeterminate moment problems*, in: *Proceedings of the International Workshop "Special Functions", Hong Kong, June 21–25, 1999*, C. F. Dunkl, M. Ismail, and R. Wong (Eds.), World Scientific, Singapore 2000, pp. 31–41.
- [4] C. Berg, *On powers of Stieltjes moment sequences. I*, J. Theoret. Probab. 18 (4) (2005), pp. 871–889.
- [5] C. Berg, *On powers of Stieltjes moment sequences. II*, J. Comput. Appl. Math. 199 (1) (2007), pp. 23–38.
- [6] C. Berg and G. Forst, *Potential Theory on Locally Compact Abelian Groups*, Ergeb. Math. Grenzgeb., Band 87, Springer, New York–Heidelberg 1975.
- [7] C. Berg and J. L. López, *Asymptotic behaviour of the Urbanik semigroup*, J. Approx. Theory 195 (2015), pp. 109–121.
- [8] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, D. Zwillinger (Ed.), eighth edition, Elsevier and Academic Press, Amsterdam 2015.
- [9] P. Hörfelt, *The moment problem for some Wiener functionals: Corrections to previous proofs*, J. Appl. Probab. 42 (3) (2005), pp. 851–860.
- [10] Z. J. Jurek, *On relations between Urbanik and Mehler semigroups*, Probab. Math. Statist. 29 (2) (2009), pp. 297–308.

- [11] G. D. Lin, *On powers of the Catalan number sequence*, Discrete Math. 342 (7) (2019), pp. 2139–2147.
- [12] G. D. Lin and J. Stoyanov, *Moment determinacy of powers and products of nonnegative random variables*, J. Theoret. Probab. 28 (4) (2015), pp. 1337–1353.
- [13] J. L. López, P. Pagola, and E. Pérez Sinusía, *A systematization of the saddle point method: Application to the Airy and Hankel functions*, J. Math. Anal. Appl. 354 (1) (2009), pp. 347–359.
- [14] F. W. J. Olver and L. C. Maximon, *Chapter 10: Bessel Functions*, in: *NIST Handbook of Mathematical Functions*, NIST and Cambridge University Press, 2010.
- [15] H. L. Pedersen, *On Krein's theorem for indeterminacy of the classical moment problem*, J. Approx. Theory 95 (1) (1998), pp. 90–100.
- [16] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions: Theory and Applications*, second edition, Walter de Gruyter & Co., Berlin 2012.
- [17] J. A. Shohat and J. D. Tamarkin, *The Problem of Moments*, American Mathematical Society, New York 1943.
- [18] A. V. Skorokhod, *Asymptotic formulas for stable distribution laws*, Dokl. Akad. Nauk SSSR (N.S.) 98 (1954), pp. 731–734.
- [19] T.-J. Stieltjes, *Recherches sur les fractions continues*, Ann. Fac. Sci. Toulouse, Math., Sér. 1, 8 (4) (1894), pp. J1–J122.
- [20] M. L. Targhetta, *On a family of indeterminate distributions*, J. Math. Anal. Appl. 147 (2) (1990), pp. 477–479.
- [21] Shu-gwei Tyan, *The structure of bivariate distribution functions and their relation to Markov processes*, Ph.D. Thesis, Princeton University, 1975.
- [22] K. Urbanik, *Functionals on transient stochastic processes with independent increments*, Studia Math. 103 (3) (1992), pp. 299–315.
- [23] V. M. Zolotarev, *One-dimensional Stable Distributions*, American Mathematical Society, Providence, RI, 1986.

Christian Berg  
Department of Mathematical Sciences  
University of Copenhagen  
Universitetsparken 5  
2100 Copenhagen Ø, Denmark  
E-mail: berg@math.ku.dk

Received on 10.2.2018;  
revised version on 29.6.2018

---